

# Week 7 Notes

Robert

July 2024

## List of Definitions

1.1	Definition (Separation) . . . . .	2
1.3	Definition (Path) . . . . .	2
1.5	Definition (Totally disconnected space) . . . . .	2

## List of Propositions

1.2	Proposition (Connected iff clopen sets are trivial) . . . . .	2
1.4	Theorem (Path connectedness implies connectedness) . . . . .	2
1.7	Theorem (Connected subspaces of $\mathbb{R}$ ) . . . . .	2
1.9	Proposition (Separations and connected subspaces) . . . . .	3
1.10	Proposition (Union of (path)-connected subspaces with a common point is (path)-connected) . . . . .	3
1.11	Proposition (Closure of a space is connected) . . . . .	3
1.12	Theorem (Main theorem on connectedness) . . . . .	3
1.15	Proposition (Finite product of connected is connected) . . . . .	4
1.16	Proposition (Product of (path) connected spaces is connected) . . . . .	4

## 1 Connectedness

**Definition 1.1 (Separation).** Let  $X$  be a topological space. Then a **separation** of  $X$  is a partition of  $X = A \cup B$ , where  $A, B$  are disjoint, open and nonempty sets.

Note that this definition may be called a *disconnection* of  $X$  by some authors (c.f. [Lee11]). A space  $X$  is **connected** if and only if there exists no separation of  $X$ .

A set is said to be **clopen** if it is both open and closed.

**Proposition 1.2** (Connected iff clopen sets are trivial). *Let  $X$  be a topological space. Then  $X$  is connected if and only if it has no nontrivial clopen subsets, i.e. the only clopen subsets of  $X$  are  $\emptyset$  and  $X$ .*

*Proof.* Obviously. □

**Definition 1.3 (Path).** Let  $X$  be a topological space and  $x, y \in X$ . Then, a **path** from  $x$  to  $y$  is a continuous function  $p : [0, 1] \rightarrow X$  such that  $p(0) = x$  and  $p(1) = y$ .

Note that the domain can be replaced with any closed interval  $[a, b]$  since all closed intervals are homeomorphic to  $[0, 1]$ .

A space  $X$  is said to be **path-connected** if given any  $x, y \in X$ , there exists a path from  $x$  to  $y$ .

**Theorem 1.4 (Path connectedness implies connectedness).** Let  $X$  be a path-connected space. Then  $X$  is connected.

*Proof.* If not, let  $A, B$  be a separation of  $X$ . Let  $a \in A, b \in B$  and  $p$  is a path from  $a$  to  $b$ . Then  $p[[0, 1]] \cap A$  and  $p[[0, 1]] \cap B$  is a separation of  $p[[0, 1]]$  which contradicts the connectedness of  $p[[0, 1]]$ . □

Note that we have made use of the fact that intervals are connected, and the image of a connected space under a continuous function is connected.

**Definition 1.5 (Totally disconnected space).** A space  $X$  is **totally disconnected** if the only connected subspaces of  $X$  are singletons.

Clearly the discrete topology on a space with more than one point is totally disconnected. However, not every totally disconnected space has the discrete topology.

**Example 1.6** (The rationals are totally disconnected). Let  $X = \mathbb{Q}$  considered as a subspace of  $\mathbb{R}$ . Then  $X$  is totally disconnected since given  $p, q$  where  $p \neq q$ , we can partition  $X = (X \cap (-\infty, p)) \cup ((p, \infty) \cap X)$ . //

Recall that if  $X \subseteq \mathbb{R}$ , a subset  $A \subseteq X$  is said to be *convex* if given  $a, b \in A$ , we have that  $[a, b] \subseteq X$ . We shall now prove that intervals in  $\mathbb{R}$  are connected. Before we begin the proof, note the properties of the real numbers that we make use of: the fact that supremums exist, and between any two reals, we can find another real.

**Theorem 1.7 (Connected subspaces of  $\mathbb{R}$ ).** Let  $X \subseteq \mathbb{R}$ . Then  $X$  is connected if and only if it is convex.

*Proof.* If  $X$  is not convex, then let  $a, b \in X$  and  $z \in \mathbb{R}$  be such that  $a < z < b$ , and  $z \notin X$ . Then  $X$  can be separated by  $X = (X \cap (-\infty, z)) \cup ((z, \infty) \cap X)$ .

Suppose  $X$  is convex but that  $X = A \cup B$  is a separation. Let  $a \in A, b \in B$  and suppose without loss of generality that  $a < b$ . Since  $X$  is convex,  $[a, b] \subseteq X$ . We thus separate  $[a, b] = (A \cap [a, b]) \cup (B \cap [a, b])$ . Let  $A_0 = (A \cap [a, b]), B_0 = (B \cap [a, b])$ . Let  $c = \sup A_0$ . Then  $c \in X$  and  $c \in A_0$  as  $A_0$  is closed. Since  $A_0$  is open there is some  $\varepsilon$  such that  $c + \varepsilon \in A_0$ . But  $c + \varepsilon > c$  which contradicts  $c$  being  $\sup A_0$ . Ooopsies! □

**Example 1.8 (Topologist's sine curve).** The topologist's sine curve is an example of a space which is connected, but not path connected. Let  $f : (0, 1] \rightarrow \mathbb{R}$  be defined by  $f(x) = \sin(1/x)$ . Let  $S \subseteq \mathbb{R}^2$  be the graph of  $f$ . The topologist's sine curve is thus defined to be  $\overline{S}$ . It is connected, because it is the closure of the image of a connected space under

a continuous function. (The function is  $x \mapsto (x, f(x))$ ). However, we run into an issue when trying to construct a path from  $x \in S$  to the set of limit points of  $S$ . For concreteness, let us suppose we are trying to connect  $x$  to  $(0, 0)$ . Suppose we somehow have a path  $p : [0, 1] \rightarrow \overline{S}$  from  $x$  to  $(0, 0)$ . Let  $L = \{0\} \times [-1, 1]$  (which is the set of limit points of  $S$ ).  $L$  is closed in  $\overline{S}$ , so  $p^{-1}(L)$  is closed too.

See [Mun00, Example 7, pp. 156–157] for full argument. (For an explicit value of  $u$  that can be chosen, you can pick  $u = \frac{1}{2n\pi + \pi/2}$  so  $\sin(1/u) = 1$ , and  $u = \frac{1}{2n\pi + (3/2)\pi}$  if you need  $\sin(1/u) = -1$ .)  $\parallel$




Figure 1: Topologist's sine curve

**Proposition 1.9** (Separations and connected subspaces). *Let  $X$  be a topological space and let  $Y$  be a (path) connected subspace of  $X$ . If  $A \cup B$  is a separation of  $X$ , then either  $Y \subseteq A$  or  $Y \subseteq B$ .*

*Proof.* If  $Y$  is not fully contained within either  $A$  or  $B$  then we can separate  $Y$  with  $(Y \cap A) \cup (Y \cap B)$ .  $\square$

**Proposition 1.10** (Union of (path)-connected subspaces with a common point is (path)-connected). *Let  $X$  be a topological space and let  $A_\alpha$  be a collection of (path) connected spaces and suppose  $z \in A_\alpha$  for all  $\alpha$ , so the  $A_\alpha$ 's have a common point. Then,  $\bigcup A_\alpha$  is (path) connected.*

*Proof.* We first prove it for path connectedness. Let  $z$  be a point in common. If  $x, y \in \bigcup A_\alpha$ , say  $x \in A_\alpha$  and  $y \in A_\beta$ . Then, glue a path from  $x$  to  $z$  and a path from  $z$  to  $y$  together. This one is easy to visualize by drawing a picture.

Let us now prove it for connectedness. Suppose that  $\bigcup A_\alpha$  is the union of disjoint open sets  $A \cup B$ . Then  $z \in A$  or  $z \in B$ . Suppose without loss of generality that  $z \in A$ . Then for all  $\alpha$ ,  $A_\alpha$  must intersect  $A$ . By the previous proposition, all  $A_\alpha \subseteq A$ . So this means  $B$  is empty. Thus there is no separation of  $\bigcup A_\alpha$ .  $\square$

**Proposition 1.11** (Closure of a space is connected). *Suppose  $A$  is a connected subspace of  $X$  and  $B$  is a set such that  $A \subseteq B \subseteq \overline{A}$ . Then,  $B$  is connected.*

*Proof.* Use Proposition 1.9. (For full proof, see [Mun00] or [Lee11, Prop 4.9, p. 88].)  $\square$

It is important to note that this proposition is untrue if  $A$  is path connected. See Example 1.8 for this happening.

**Theorem 1.12 (Main theorem on connectedness).** *Let  $X$  be a connected space and let  $f : X \rightarrow Y$  be a continuous function. Then  $f[X]$  is connected.*

*Proof.* If not, let  $A, B$  be a separation of  $f[X]$ . Then  $f^{-1}(A), f^{-1}(B)$  separate  $X$ .  $\square$

Note here that  $A, B$  are considered as open/closed sets in the subspace topology on  $f[X]$ . The above theorem is also true with path-connectedness in place of connectedness. The proof is obvious, as you can simply compose the path with  $f$ .

**Corollary 1.13** (Connectedness is invariant under homeomorphism). *Any space homeomorphic to a connected space is connected.*

*Proof.* Duh. □

**Corollary 1.14** (Intermediate value theorem). *Let  $f : X \rightarrow \mathbb{R}$  and suppose  $X$  is connected. If  $p, q \in X$  then  $f$  attains every value between  $f(p)$  and  $f(q)$ .*

*Proof.* Suppose without loss of generality that  $f(p) < f(q)$ . Then  $f[X]$  is connected so it must contain  $[f(p), f(q)]$ . □

See [Lee11, Thm 4.12, p. 89] for further details.

**Warning.** The preimage of a connected or path-connected space need not be connected. Take  $X = \mathbb{R}$  with the discrete topology and  $Y = \mathbb{R}$  with the trivial topology. Then the identity is continuous from  $X$  to  $Y$  but the preimage of  $Y$  is disconnected.

**Proposition 1.15** (Finite product of connected is connected). *If  $X, Y$  are connected spaces then  $X \times Y$  is connected.*




Figure 2: Proof that the finite product of connected spaces is connected

*Proof.* Fix a point  $\langle a, b \rangle \in X \times Y$ . Define  $T_x = \{x\} \times Y \cup X \times \{b\}$ . This set is connected as it is the union of 2 connected sets with the point  $\langle a, b \rangle$  in common. Then  $X \times Y = \bigcup_{x \in X} T_x$ . This is a union of connected spaces with the point  $\langle a, b \rangle$  in common. (See [Section 1](#) for a better visualization. See [Mun00, Thm 23.6, p. 148] for complete proof.) □

The product topology preserves connectedness (which is nice).

**Proposition 1.16** (Product of (path) connected spaces is connected). *If  $X_\alpha$  is a collection of (path)-connected spaces, then  $X = \prod X_\alpha$  is (path)-connected.*

*Proof.* (Path-connectedness) Let  $\mathbf{x}, \mathbf{y} \in \prod X_\alpha$ , writing  $\mathbf{x} = \langle x_\alpha : \alpha \in \Lambda \rangle$  and  $\mathbf{y} = \langle y_\alpha : \alpha \in \Lambda \rangle$ . Since each  $X_\alpha$  is path connected, for each  $\alpha$ , let  $f_\alpha : I \rightarrow X_\alpha$  be a path from  $x_\alpha$  to  $y_\alpha$ . We simply glue these paths together by taking  $f(t) = \langle f_\alpha(t) : \alpha \in \Lambda \rangle$  which is a path from  $\mathbf{x}$  to  $\mathbf{y}$ .

(Connectedness) Fix a point  $\mathbf{a} = \langle a_\alpha : \alpha \in \Lambda \rangle$ . If  $F \subseteq \Lambda$  is finite, then define

$$X_F = \{ \mathbf{x} \in X : x_\alpha = a_\alpha \text{ if } \alpha \in \Lambda \setminus F \}.$$

So  $X_F$  is the set of all  $\mathbf{x} \in X$  such that  $x_\alpha = a_\alpha$  for all coordinates except those in  $F$ . We thus see that  $X_F$  is homeomorphic to  $\prod_{\alpha \in F} X_\alpha$ . Since finite products of connected spaces are connected,  $X_F$  is connected.

Now, set  $Z = \bigcup_{F \subseteq \Lambda, |F| < \omega} X_F$ . This is the union of  $X_F$ 's across all finite subsets  $F \subseteq \Lambda$ . Then  $Z$  is connected, as each  $X_F$  has the point  $\mathbf{a}$  in common ([Proposition 1.10](#)). Additionally, we claim that  $\overline{Z} = X$ . This will finish it off ([Proposition 1.11](#)), so let us see why this is true. Pick  $\mathbf{x} \in X$  and let  $U$  be a neighborhood of  $\mathbf{x}$  in the product topology. We need to show that  $U$  intersects  $Z$ . Since we are in the product topology, this means that  $U = \prod_{\alpha \in \Lambda} U_\alpha$  and  $U_\alpha = X_\alpha$  except for finite  $\alpha$ . Say those  $\alpha$ 's are all in the set  $F$ . Define

$$\mathbf{z}_\alpha = \begin{cases} x_\alpha & \text{if } \alpha \in F, \\ a_\alpha & \text{otherwise.} \end{cases}$$

Then  $\mathbf{z} \in X_F \cap U$  so the point  $\mathbf{z}$  is in the closure of  $Z$ . □

The box topology is usually not going to be connected.

**Example 1.17** (Countable product of  $\mathbb{R}$  with the box topology). Let  $X = \prod_{n \in \mathbb{N}} \mathbb{R}$  and give it the box topology. Let

$$\ell^\infty = \{ \mathbf{x} \in \mathbb{R}^{\mathbb{N}} : \sup_{n \in \mathbb{N}} |x_n| < \infty \}.$$

This is the set of bounded real-valued sequences. We shall show that  $\ell^\infty$  is clopen. Since  $\ell^\infty$  is not all of  $\mathbb{R}^{\mathbb{N}}$  (by obviousness) we will be done. Let  $\mathbf{x} \in \ell^\infty$  be a bounded sequence. Consider the neighborhood of  $\mathbf{x}$  given by  $U = \prod_{n \in \mathbb{N}} B(x_n, 1)$ . Notice if  $\mathbf{y} \in U$ , then  $|y_n| < |x_n| + 1 \leq \sup_{n \in \mathbb{N}} |x_n| + 1$  so  $\mathbf{y}$  must be a bounded sequence too. For being closed, notice that the complement is open. (Use the same argument). //

**Remark 1.18** (Path-connectedness of finite products). For the finite case the proof is very easy. Given a point  $\langle x_0, y_0 \rangle \in X \times Y$  and a point  $\langle x_1, y_1 \rangle \in X \times Y$ , since  $X, Y$  are respectively path connected let  $p$  be a path in  $X$  from  $x_0$  to  $x_1$  and  $q$  be a path in  $Y$  from  $y_0$  to  $y_1$ . Then the map  $p \times q$  is the desired path. Apply induction and the fact that  $(X \times Y) \times Z$  is homeomorphic to  $X \times Y \times Z$ .